arXiv:math-ph/9902017v1 12 Feb 1999

Feb. 9, 1999
PUPT-1832

On Lieb's conjecture for the Wehrl entropy of Bloch coherent states

Peter Schupp

Department of Physics, Princeton University
Princeton, NJ 08544-0708, USA

Abstract

Lieb's conjecture for the Wehrl entropy of Bloch coherent states is proved for spin 1 and spin 3/2. Using a geometric representation we solve the entropy integrals for states of arbitrary spin and evaluate them explicitly in the cases of spin 1, 3/2, and 2. We also give a group theoretic proof for all spin of a related inequality.

©1999 by the author. Reproduction of this article, in its entirety, by any means is permitted for non-commercial purposes.

e-mail: schupp@princeton.edu

1 Introduction

Wehrl proposed [16] a hybrid between quantum mechanical and classical entropy that enjoys monotonicity, strong subadditivity and positivity – physically desirable properties, some of which both kinds of entropy lack [7, 8]. This new entropy is the ordinary Shannon entropy of the probability density provided by the lower symbol of the density matrix.

For a quantum mechanical system with density matrix ρ, Hilbert space $\mathcal{H}$, and a family of normalized coherent states $|z\rangle$, parametrized symbolically by z and satisfying $\int dz\, |z\rangle\langle z| = \mathbf{1}$ (resolution of identity), the Wehrl entropy is

$$S_W(\rho) = -\int dz\, \langle z|\rho|z\rangle \ln\langle z|\rho|z\rangle. \tag{1}$$

Like quantum mechanical entropy, $S_Q = -\mathrm{tr}\, \rho \ln \rho$, Wehrl entropy is always non-negative, in fact $S_W > S_Q \geq 0$. In view of this inequality it is interesting to ask for the minimum of S_W and the corresponding minimizing density matrix. It follows from concavity of $-x \ln x$ that a minimizing density matrix must be a pure state, *i.e.*, $\rho = |\psi\rangle\langle\psi|$ for a normalized vector $|\psi\rangle \in \mathcal{H}$ [6]. (Note that $S_W(|\psi\rangle\langle\psi|)$ depends on $|\psi\rangle$ and is non-zero, unlike the quantum entropy which is of course zero for pure states.)

For Glauber coherent states Wehrl conjectured [16] and Lieb proved [6] that the minimizing state $|\psi\rangle$ is again a coherent state. It turns out that all Glauber coherent states have Wehrl entropy one, so Wehrl's conjecture can be written as follows:

THEOREM 1.1 (Lieb) *The minimum of $S_W(\rho)$ for states in $\mathcal{H} = L^2(\mathbb{R})$ is one,*

$$S_W(|\psi\rangle\langle\psi|) = -\int dz\, |\langle\psi|z\rangle|^2 \ln |\langle\psi|z\rangle|^2 \geq 1, \tag{2}$$

with equality if and only if $|\psi\rangle$ is a coherent state.

To prove this, Lieb used a clever combination of the sharp Hausdorff-Young inequality [2, 9, 11] and the sharp Young inequality [3, 2, 9, 11] to show that

$$s \int dz \, |\langle z|\psi\rangle|^{2s} \leq 1, \quad s \geq 1, \tag{3}$$

again with equality if and only if $|\psi\rangle$ is a coherent state. Wehrl's conjecture follows from this in the limit $s \to 1$ essentially because (2) is the derivative of (3) with respect to s at $s = 1$. All this easily generalizes to $L^2(\mathbb{R}^n)$ [6, 10].

The lower bound on the Wehrl entropy is related to Heisenberg's uncertainty principle [1, 4] and it has been speculated that S_W can be used to measure uncertainty due to both quantum and thermal fluctuations [4].

It is very surprising that 'heavy artillery' like the sharp constants in the mentioned inequalities are needed in Lieb's proof. To elucidate this situation, Lieb suggested [6] studying the analog of Wehrl's conjecture for Bloch coherent states $|\Omega\rangle$, where one should expect significant simplification since these are finite dimensional Hilbert spaces. However, no progress has been made, not even for a single spin, even though many attempts have been made [7]. Attempts to proceed again along the lines of Lieb's original proof have failed to provide a sharp inequality and the direct computation of the entropy and related integrals, even numerically, was unsuccesful [14].

The key to the recent progress is a geometric representation of a state of spin j as $2j$ points on a sphere. In this representation the expression $|\langle\Omega|\psi\rangle|^2$ factorizes into a product of $2j$ functions f_i on the sphere, which measure the square chordal distance from the antipode of the point parametrized by Ω to each of the $2j$ points on the sphere. Lieb's conjecture, in a generalized form analogous to (3), then looks like the quotient of two Hölder inequalities

$$\frac{\|f_1 \cdots f_{2j}\|_s}{\|f_1 \cdots f_{2j}\|_1} \leq \frac{\prod_{i=1}^{2j} \|f_i\|_{2js}}{\prod_{i=1}^{2j} \|f_i\|_{2j}}, \tag{4}$$

with the one with the higher power winning against the other one. We shall give a group theoretic proof of this inequality for the special case $s \in \mathbb{N}$ in theorem 4.3.

In the geometric representation the Wehrl entropy of spin states finds a direct physical interpretation: It is the classical entropy of a single particle on a sphere interacting via Coulomb potential with $2j$ fixed sources; s plays the role of inverse temperature.

The entropy integral (1) can now be done because $|\langle\Omega|\psi\rangle|^2$ factorizes and one finds a formula for the Wehrl entropy of any state. When we evaluate the entropy explicitly for states of spin 1, 3/2, and 2 we find surprisingly simple expressions solely in terms of the square chordal distances between the points on the sphere that define the given state.

A different, more group theoretic approach seems to point to a connection between Lieb's conjecture and the norm of certain spin js states with $1 \leq s \in \mathbb{R}$ [5]. So far, however, this has only been useful for proving the analog of inequality (3) for $s \in \mathbb{N}$.

We find that a proof of Lieb's conjecture for low spins can be reduced to some beautiful spherical geometry, but the unreasonable difficulty of a complete proof is still a great puzzle; its resolution may very well lead to interesting mathematics and perhaps physics.

2 Bloch coherent spin states

Glauber coherent states $|z\rangle = \pi^{-\frac{1}{4}} e^{-(x-q)^2/2} e^{ipx}$, parametrized by $z = (q + ip)/\sqrt{2}$ and with measure $dz = dpdq/2\pi$, are usually introduced as eigenvectors of the annihilation operator $a = (\hat{x} + i\hat{p})/\sqrt{2}$, $a|z\rangle = z|z\rangle$, but the same states can also be obtained by the action of the Heisenberg-Weyl group $H_4 = \{a^\dagger a, a^\dagger, a, I\}$ on the extremal state $|0\rangle = \pi^{-\frac{1}{4}} e^{-x^2/2}$. Glauber coherent states are thus elements of the coset space of the Heisenberg-Weyl group modulo the stability subgroup $U(1) \otimes U(1)$ that leaves the extremal state invariant. (See *e.g.* [17] and references therein.) This construction easily generalizes to other groups, in particular to SU(2), where it gives the Bloch coherent spin states [13] that we are interested in: Here the Hilbert space can be any one of the finite dimensional spin-j representations $[j] \equiv \mathbb{C}^{2j+1}$ of SU(2), $j = \frac{1}{2}, 1, \frac{3}{2}, \ldots$, and the extremal state for each $[j]$ is the highest weight vector $|j,j\rangle$. The stability subgroup is U(1) and the coherent states are thus elements of the sphere S_2 =SU(2)/U(1); they can be labeled by $\Omega = (\theta, \phi)$ and are obtained from $|j,j\rangle$ by rotation:

$$|\Omega\rangle_j = \mathcal{R}_j(\Omega)|j,j\rangle. \tag{5}$$

For spin $j = \frac{1}{2}$ we find

$$|\omega\rangle = p^{\frac{1}{2}} e^{-i\frac{\phi}{2}} |\!\uparrow\rangle + (1-p)^{\frac{1}{2}} e^{i\frac{\phi}{2}} |\!\downarrow\rangle, \tag{6}$$

with $p \equiv \cos^2\frac{\theta}{2}$. (Here and in the following $|\omega\rangle$ is short for the spin-$\frac{1}{2}$ coherent state $|\Omega\rangle_{\frac{1}{2}}$; $\omega = \Omega = (\theta, \phi)$. $|\!\uparrow\rangle \equiv |\frac{1}{2}, \frac{1}{2}\rangle$ and $|\!\downarrow\rangle \equiv |\frac{1}{2}, -\frac{1}{2}\rangle$.) An important observation for what follows is that the product of two coherent states for the same Ω is again a coherent state:

$$\begin{aligned} |\Omega\rangle_j \otimes |\Omega\rangle_{j'} &= (\mathcal{R}_j \otimes \mathcal{R}_{j'}) \left(|j,j\rangle \otimes |j',j'\rangle\right) \\ &= \mathcal{R}_{j+j'} \, |j+j', j+j'\rangle \;=\; |\Omega\rangle_{j+j'}. \end{aligned} \tag{7}$$

Coherent states are in fact the only states for which the product of a spin-j state with a spin-j' state is a spin-$(j+j')$ state and not a more general element of $[j+j'] \oplus \ldots \oplus [\,|j-j'|\,]$. From this key property an explicit representation for Bloch coherent states of higher spin can be easily derived:

$$\begin{aligned} |\Omega\rangle_j &= (|\omega\rangle)^{\otimes 2j} \;=\; \left(p^{\frac{1}{2}} e^{-i\frac{\phi}{2}} |\!\uparrow\rangle + (1-p)^{\frac{1}{2}} e^{i\frac{\phi}{2}} |\!\downarrow\rangle\right)^{\otimes 2j} \\ &= \sum_{m=-j}^{j} \binom{2j}{j+m}^{\frac{1}{2}} p^{\frac{j+m}{2}} (1-p)^{\frac{j-m}{2}} e^{-im\frac{\phi}{2}} |j,m\rangle. \end{aligned} \tag{8}$$

(The same expression can also be obtained directly from (5), see *e.g.* [12, chapter 4].) The coherent states as given are normalized $\langle\Omega|\Omega\rangle_j = 1$ and satisfy

$$(2j+1) \int \frac{d\Omega}{4\pi} \, |\Omega\rangle_j \langle\Omega|_j = P_j, \qquad \text{(resolution of identity)} \tag{9}$$

where $P_j = \sum |j,m\rangle\langle j,m|$ is the projector onto $[j]$. It is not hard to compute the Wehrl entropy for a coherent state $|\Omega'\rangle$: Since the integral over the sphere is invariant under rotations it is enough to consider the coherent state $|j,j\rangle$; then use $|\langle j,j|\Omega\rangle|^2 = |\langle\uparrow|\omega\rangle|^{2\cdot 2j} = p^{2j}$ and $d\Omega/4\pi = -dp\,d\phi/2\pi$, where $p = \cos^2\frac{\theta}{2}$ as above, to obtain

$$\begin{aligned} S_W(|\Omega'\rangle\langle\Omega'|) &= -(2j+1)\int\frac{d\Omega}{4\pi}\,|\langle\Omega|\Omega'\rangle|^2\ln|\langle\Omega|\Omega'\rangle|^2 \\ &= -(2j+1)\int_0^1 dp\,p^{2j}\,2j\ln p \;=\; \frac{2j}{2j+1}. \end{aligned} \tag{10}$$

Similarly, for later use,

$$(2js+1)\int\frac{d\Omega}{4\pi}|\langle\Omega'|\Omega\rangle|^{2s} = (2js+1)\int_0^1 dp\,p^{2js} = 1. \tag{11}$$

As before the density matrix that minimizes S_W must be a pure state $|\psi\rangle\langle\psi|$. The analog of theorem 1.1 for spin states is:

Conjecture 2.1 (Lieb) *The minimum of S_W for states in $\mathcal{H} = \mathbb{C}^{2j+1}$ is $2j/(2j+1)$,*

$$S_W(|\psi\rangle\langle\psi|) = -(2j+1)\int\frac{d\Omega}{4\pi}\,|\langle\Omega|\psi\rangle|^2\ln|\langle\Omega|\psi\rangle|^2 \geq \frac{2j}{2j+1}, \tag{12}$$

with equality if and only if $|\psi\rangle$ is a coherent state.

Remark: For spin 1/2 this is an identity because all spin 1/2 states are coherent states. The first non-trivial case is spin $j = 1$.

3 Proof of Lieb's conjecture for low spin

In this section we shall geometrize the description of spin states, use this to solve the entropy integrals for all spin and prove Lieb's conjecture for low spin by actual computation of the entropy.

LEMMA 3.1 *States of spin j are in one to one correspondence to $2j$ points on the sphere S_2: With $2j$ points, parametrized by $\omega_k = (\theta_k, \phi_k)$, $k = 1, \ldots, 2j$, we can associate a state*

$$|\psi\rangle = c^{\frac{1}{2}} P_j(|\omega_1\rangle \otimes \ldots \otimes |\omega_{2j}\rangle) \ \in \ [j], \tag{13}$$

and every state $|\psi\rangle \in [j]$ is of that form. (The spin-$\frac{1}{2}$ states $|\omega_k\rangle$ are given by (6), $c^{\frac{1}{2}} \neq 0$ fixes the normalization of $|\psi\rangle$, and P_j is the projector onto spin j.)

Remark: Some or all of the points may coincide. Coherent states are exactly those states for which all points on the sphere coincide. $c^{\frac{1}{2}} \in \mathbb{C}$ may contain an (unimportant) phase that we can safely ignore in the following. This representation is unique up to permutation of the $|\omega_k\rangle$. The ω_k may be found by looking at $\langle\Omega|\psi\rangle$ as a function of $\Omega = (\theta, \psi)$: they are the antipodal points to the zeroes of this function.

PROOF: Rewrite (8) in complex coordinates for $\theta \neq 0$

$$z = \left(\frac{p}{1-p}\right)^{\frac{1}{2}} e^{i\phi} = \cot\frac{\theta}{2} e^{i\phi} \tag{14}$$

(stereographic projection) and contract it with $|\psi\rangle$ to find

$$\langle\Omega|\psi\rangle = \frac{e^{-ij\phi}}{(1+z\bar{z})^j} \sum_{m=-j}^{j_{\max}} \begin{pmatrix} 2j \\ j+m \end{pmatrix}^{\frac{1}{2}} z^{j+m} \psi_m, \tag{15}$$

where $j_{\max}$ is the largest value of m for which ψ_m in the expansion $|\psi\rangle = \sum \psi_m |m\rangle$ is nonzero. This is a polynomial of degree $j + j_{\max}$ in $z \in \mathbb{C}$ and can thus be factorized:

$$\langle\Omega|\psi\rangle = \frac{e^{-ij\phi}\psi_{j_{\max}}}{(1+z\bar{z})^j} \prod_{k=1}^{j+j_{\max}} (z - z_k). \tag{16}$$

Consider now the spin $\frac{1}{2}$ states $|\omega_k\rangle = (1 + z_k\bar{z}_k)^{-\frac{1}{2}}(|\uparrow\rangle - z_k|\downarrow\rangle)$ for $1 \leq k \leq j + j_{\max}$ and $|\omega_m\rangle = |\downarrow\rangle$ for $j + j_{\max} < m \leq 2j$. According to (15):

$$\langle\omega|\omega_k\rangle = \frac{e^{-\frac{i\phi}{2}}}{(1+z\bar{z})^{\frac{1}{2}}(1+z_k\bar{z}_k)^{\frac{1}{2}}}(z - z_k), \qquad \langle\omega|\omega_m\rangle = \frac{e^{-\frac{i\phi}{2}}}{(1+z\bar{z})^{\frac{1}{2}}}, \tag{17}$$

so by comparison with (16) and with an appropriate constant c

$$\langle\Omega|\psi\rangle = c^{\frac{1}{2}}\langle\omega|\omega_1\rangle\cdots\langle\omega|\omega_{2j}\rangle = c^{\frac{1}{2}}\langle\Omega|\omega_1\otimes\ldots\otimes\omega_{2j}\rangle. \tag{18}$$

By inspection we see that this expression is still valid when $\theta = 0$ and with the help of (9) we can complete the proof the lemma. ■

We see that the geometric representation of spin states leads to a factorization of $\langle\Omega|\psi\rangle|^2$. In this representation we can now do the entropy integrals, essentially because the logarithm becomes a simple sum.

THEOREM 3.2 *Consider any state $|\psi\rangle$ of spin j. According to lemma 3.1, it can be written as $|\psi\rangle = c^{\frac{1}{2}}P_j(|\omega_1\rangle\otimes\ldots\otimes|\omega_{2j}\rangle)$. Let $\mathcal{R}_i$ be the rotation that turns ω_i to the 'north pole', $\mathcal{R}_i|\omega_i\rangle = |\uparrow\rangle$, let $|\psi^{(i)}\rangle = \mathcal{R}_i|\psi\rangle$, and let $\psi_m^{(i)}$ be the coefficient of $|j,m\rangle$ in the expansion of $|\psi^{(i)}\rangle$, then the Wehrl entropy is:*

$$S_W(|\psi\rangle\langle\psi|) = \sum_{i=1}^{2j}\sum_{m=-j}^{j}\left(\sum_{n=0}^{j-m}\frac{1}{2j+1-n}\right)|\psi_m^{(i)}|^2 - \ln c. \tag{19}$$

Remark: This formula reduces the computation of the Wehrl entropy of any spin state to its factorization in the sense of lemma 3.1, which in general requires the solution of a $2j$'th order algebraic equation. This may explain why previous attempts to do the entropy integrals have failed. The $n = 0$ terms in the expression for the entropy sum up to $2j/(2j+1)$, the entropy of a coherent state, and Lieb's conjecture can be thus be written

$$\ln c \leq \sum_{i=1}^{2j}\sum_{m=-j+1}^{j-1}\left(\sum_{n=1}^{j-m}\frac{1}{2j+1-n}\right)|\psi_m^{(i)}|^2. \tag{20}$$

Note that $\psi_{-j}^{(i)} = 0$ by construction of $|\psi^{(i)}\rangle$: $\psi_{-j}^{(i)}$ contains a factor $\langle\downarrow|\uparrow\rangle$.
A similar calculation gives

$$\ln c = 2j + \int\frac{d\Omega}{4\pi}\ln|\langle\Omega|\psi\rangle|^2. \tag{21}$$

PROOF: Using lemma 3.1, (9), the rotational invariance of the measure and the inverse Fourier transform in ϕ we find

$$\begin{aligned} S_W(|\psi\rangle\langle\psi|) &= -(2j+1)\int \frac{d\Omega}{4\pi}|\langle\Omega|\psi\rangle|^2 \sum_{i=1}^{2j} \ln|\langle\omega|\omega_i\rangle|^2 - \ln c \\ &= -(2j+1)\sum_{i=1}^{2j}\int \frac{d\Omega}{4\pi}|\langle\Omega|\psi^{(i)}\rangle|^2 \ln|\langle\omega|\uparrow\rangle|^2 - \ln c \\ &= -(2j+1)\sum_{i=1}^{2j}\sum_{m=-j}^{j} |\psi^{(i)}_m|^2 \tbinom{2j}{j+m} \int_0^1 dp\, p^{j+m}(1-p)^{j-m}\ln p - \ln c. \end{aligned} \tag{22}$$

It is now easy to do the remaining p-integral by partial integration to proof the theorem. ■

Lieb's conjecture for low spin can be proved with the help of formula (19). For spin 1/2 there is nothing to prove, since all states of spin 1/2 are coherent states. The first nontrivial case is spin 1:

COROLLARY 3.3 (spin 1) *Consider an arbitrary state of spin 1. Let* μ *be the square of the chordal distance between the two points on the sphere of radius* $\frac{1}{2}$ *that represent this state. It's Wehrl entropy is given by*

$$S_W(\mu) = \frac{2}{3} + c\cdot\left(\frac{\mu}{2} + \frac{1}{c}\ln\frac{1}{c}\right), \tag{23}$$

with

$$\frac{1}{c} = 1 - \frac{\mu}{2}. \tag{24}$$

Lieb's conjecture holds for all states of spin 1: $S_W(\mu) \geq 2/3 = 2j/(2j+1)$ *with equality for* $\mu = 0$, i.e. *for coherent states.*

PROOF: Because of rotational invariance we can assume without loss of generality that the first point is at the 'north pole' of the sphere and that the second point is parametrized as $\omega_2 = (\tilde{\theta}, \tilde{\phi} = 0)$, so that $\mu = \sin^2 \frac{\tilde{\theta}}{2}$. Up to normalization (and an irrelevant phase)

$$|\tilde{\psi}\rangle = P_{j=1} |\uparrow \otimes \tilde{\omega}\rangle \tag{25}$$

is the state of interest. But from (6)

$$|\uparrow \otimes \tilde{\omega}\rangle = (1-\mu)^{\frac{1}{2}} |\uparrow\uparrow\rangle + \mu^{\frac{1}{2}} |\uparrow\downarrow\rangle. \tag{26}$$

Projecting onto spin 1 and inserting the normalization constant $c^{\frac{1}{2}}$ we find

$$|\psi\rangle = c^{\frac{1}{2}} \left((1-\mu)^{\frac{1}{2}} |1,1\rangle + \mu^{\frac{1}{2}} \frac{1}{\sqrt{2}} |1,0\rangle \right). \tag{27}$$

This gives (ignoring a possible phase)

$$1 = \langle\psi|\psi\rangle = c\left(1 - \mu + \frac{\mu}{2}\right) = c\left(1 - \frac{\mu}{2}\right) \tag{28}$$

and so $1/c = 1 - \mu/2$. Now we need to compute the components of $|\psi^{(1)}\rangle$ and $|\psi^{(2)}\rangle$. Note that $|\psi^{(1)}\rangle = |\psi\rangle$ because ω_1 is already pointing to the 'north pole'. To obtain $|\psi^{(2)}\rangle$ we need to rotate point 2 to the 'north pole'. We can use the remaining rotational freedom to effectively exchange the two points, thereby recovering the original state $|\psi\rangle$. The components of both $|\psi^{(1)}\rangle$ and $|\psi^{(2)}\rangle$ can thus be read off (27):

$$\psi_1^{(1)} = \psi_1^{(2)} = c^{\frac{1}{2}} (1-\mu)^{\frac{1}{2}}, \qquad \psi_0^{(1)} = \psi_0^{(2)} = c^{\frac{1}{2}} \mu^{\frac{1}{2}} / \sqrt{2}. \tag{29}$$

Inserting now c, $|\psi_1^{(1)}|^2 = |\psi_1^{(2)}|^2 = c(1-\mu)$, and $|\psi_0^{(1)}|^2 = |\psi_0^{(2)}|^2 = c\mu/2$ into (19) gives the stated entropy.

To prove Lieb's conjecture for states of spin 1 we use (28) to show that the second term in (23) is always non-negative and zero only for $\mu = 0$, *i.e.* for a coherent state. This follows from

$$\frac{c\mu}{2} - \ln c \geq \frac{c\mu}{2} + 1 - c = 0 \tag{30}$$

with equality for $c = 1$ which is equivalent to $\mu = 0$. ■



Figure 1: Spin 3/2

Corollary 3.4 (spin 3/2) *Consider an arbitrary state of spin 3/2. Let* ϵ, μ, ν *be the squares of the chordal distances between the three points on the sphere of radius* $\frac{1}{2}$ *that represent this state (see figure 1). It's Wehrl entropy is given by*

$$S_W(\epsilon,\mu,\nu)=\frac{3}{4}+c\cdot\left(\frac{\epsilon+\mu+\nu}{3}-\frac{\epsilon\mu+\epsilon\nu+\mu\nu}{6}+\frac{1}{c}\ln\frac{1}{c}\right) \tag{31}$$

with

$$\frac{1}{c}=1-\frac{\epsilon+\mu+\nu}{3}. \tag{32}$$

Lieb's conjecture holds for all states of spin 3/2: $S_W(\epsilon,\mu,\nu)\geq 3/4=2j/(2j+1)$ *with equality for* $\epsilon=\mu=\nu=0$, i.e. *for coherent states.*

Proof: The proof is similar to the spin 1 case, but the geometry and algebra is more involved. Consider a sphere of radius $\frac{1}{2}$, with points 1, 2, 3 on its surface, and two planes through its center; the first plane containing points 1 and 3, the second plane containing points 2 and 3. The intersection angle ϕ of these two planes satisfies

$$2\cos\phi\sqrt{\epsilon\mu(1-\epsilon)(1-\mu)}=\epsilon+\mu-\nu-2\epsilon\mu. \tag{33}$$

ϕ is the azimuthal angle of point 2, if point 3 is at the 'north pole' of the sphere and point 1 is assigned zero azimuthal angle.

The states $|\psi^{(1)}\rangle$, $|\psi^{(2)}\rangle$, and $|\psi^{(3)}\rangle$ all have one point at the north pole of the sphere. It is enough to compute the values of $|\psi_m^{(i)}|^2$ for one i, the other values can be found by appropriate permutation of ϵ, μ, ν. (Note that we make no restriction on the parameters $0 \leq \epsilon$, μ, $\nu \leq 1$ other than that they are square chordal distances between three points on a sphere of radius $\frac{1}{2}$.) We shall start with $i = 3$: Without loss of generality the three points can be parametrized as $\omega_1^{(3)} = (\tilde{\theta}, 0)$, $\omega_2^{(3)} = (\theta, \phi)$, and $\omega_3^{(3)} = (0, 0)$ with $\mu = \sin^2 \frac{\tilde{\theta}}{2}$ and $\epsilon = \sin^2 \frac{\theta}{2}$. Corresponding spin-$\frac{1}{2}$ states are

$$
\begin{aligned}
|\omega_1^{(3)}\rangle &= (1-\mu)^{\frac{1}{2}}|\uparrow\rangle + \mu^{\frac{1}{2}}|\downarrow\rangle, && (34)\\
|\omega_2^{(3)}\rangle &= (1-\epsilon)^{\frac{1}{2}}e^{\frac{-i\phi}{2}}|\uparrow\rangle + \epsilon^{\frac{1}{2}}e^{\frac{i\phi}{2}}|\downarrow\rangle, && (35)\\
|\omega_3^{(3)}\rangle &= |\uparrow\rangle, && (36)
\end{aligned}
$$

and up to normalization, the state of interest is

$$
\begin{aligned}
|\tilde{\psi}^{(3)}\rangle &= P_{j=3/2}|\omega_1^{(3)} \otimes \omega_2^{(3)} \otimes \omega_3^{(3)}\rangle \\
&= (1-\epsilon)^{\frac{1}{2}}(1-\mu)^{\frac{1}{2}}e^{\frac{-i\phi}{2}}|\frac{3}{2}, \frac{3}{2}\rangle \\
&\quad + \left((1-\mu)^{\frac{1}{2}}\epsilon^{\frac{1}{2}}e^{\frac{i\phi}{2}} + \mu^{\frac{1}{2}}(1-\epsilon)^{\frac{1}{2}}e^{\frac{-i\phi}{2}}\right)\frac{1}{\sqrt{3}}|\frac{3}{2}, \frac{1}{2}\rangle \\
&\quad + \mu^{\frac{1}{2}}\epsilon^{\frac{1}{2}}e^{\frac{i\phi}{2}}\frac{1}{\sqrt{3}}|\frac{3}{2}, -\frac{1}{2}\rangle. && (37)
\end{aligned}
$$

This gives

$$
\begin{aligned}
|\tilde{\psi}^{(3)}_{\frac{3}{2}}|^2 &= (1-\epsilon)(1-\mu), && (38)\\
|\tilde{\psi}^{(3)}_{\frac{1}{2}}|^2 &= \frac{1}{3}\left(\epsilon(1-\mu) + \mu(1-\epsilon) + 2\sqrt{\epsilon\mu(1-\mu)(1-\epsilon)}\cos\phi\right) \\
&= \frac{2}{3}\epsilon(1-\mu) + \frac{2}{3}\mu(1-\epsilon) - \frac{\nu}{3}, && (39)\\
|\tilde{\psi}^{(3)}_{-\frac{1}{2}}|^2 &= \frac{\epsilon\mu}{3}, && (40)
\end{aligned}
$$

and $|\tilde{\psi}^{(3)}_{-\frac{3}{2}}|^2 = 0$. The sum of these expressions is

$$
\frac{1}{c} = \langle\tilde{\psi}|\tilde{\psi}\rangle = 1 - \frac{\epsilon + \mu + \nu}{3}, \qquad (41)
$$



Figure 2: Spin 2

with $0 < 1/c \leq 1$. The case $i = 1$ is found by exchanging $\mu \leftrightarrow \nu$ (and also $3 \leftrightarrow 1$, $\phi \leftrightarrow -\phi$). The case $i = 2$ is found by permuting $\epsilon \to \mu \to \nu \to \epsilon$ (and also $1 \to 3 \to 2 \to 1$). Using (19) then gives the stated entropy.

To complete the proof Lieb's conjecture for all states of spin 3/2 we need to show that the second term in (31) is always non-negative and zero only for $\epsilon = \mu = \nu = 0$. From the inequality $(1-x)\ln(1-x) \geq -x + x^2/2$ for $0 \leq x < 1$, we find

$$\frac{1}{c}\ln\frac{1}{c} \geq -\frac{\epsilon+\mu+\nu}{3} + \frac{1}{2}\left(\frac{\epsilon+\mu+\nu}{3}\right)^2, \tag{42}$$

with equality for $c = 1$. Using the inequality between algebraic and geometric mean it is not hard to see that

$$\left(\frac{\epsilon+\mu+\nu}{3}\right)^2 \geq \frac{\epsilon\mu+\nu\epsilon+\mu\nu}{3} \tag{43}$$

with equality for $\epsilon = \mu = \nu$. Putting everything together and inserting it into (31) we have, as desired, $S_W \geq 3/4$ with equality for $\epsilon = \mu = \nu = 0$, *i.e.* for coherent states. ■

COROLLARY 3.5 (spin 2) *Consider an arbitrary state of spin 2. Let* ϵ, μ, ν, α, β, γ *be the squares of the chordal distances between the four points on the* *sphere of radius $\frac{1}{2}$ that represent this state (see figure). It's Wehrl entropy is given by*

$$S_W(\epsilon, \mu, \nu, \alpha, \beta) = \frac{4}{5} + c \cdot \left(\sigma + \frac{1}{c} \ln \frac{1}{c}\right), \tag{44}$$

where

$$\frac{1}{c} = 1 - \frac{1}{4} \sum \textcircled{|} + \frac{1}{12} \sum \textcircled{||} \tag{45}$$

and

$$\sigma = \frac{1}{12}\left(-\frac{1}{2}\sum \textcircled{\top} - \frac{5}{3}\sum \textcircled{||} - \sum \textcircled{\ulcorner} + 3\sum \textcircled{|}\right) \tag{46}$$

with

$$\sum \textcircled{\top} \equiv \alpha\mu\nu + \epsilon\beta\nu + \epsilon\mu\gamma + \alpha\beta\gamma, \tag{47}$$

$$\sum \textcircled{||} \equiv \alpha\epsilon + \beta\mu + \gamma\nu, \qquad \sum \textcircled{|} \equiv \alpha + \beta + \gamma + \mu + \nu + \epsilon, \tag{48}$$

$$\sum \textcircled{\ulcorner} \equiv \alpha\mu+\alpha\nu+\mu\nu+\beta\epsilon+\beta\nu+\epsilon\nu+\epsilon\gamma+\epsilon\mu+\mu\gamma+\alpha\beta+\alpha\gamma+\beta\gamma. \tag{49}$$

Remark: The fact that the four points lie on the surface of a sphere imposes a complicated constraint on the parameters ϵ, μ, ν, α, β, γ. Although we have convincing numerical evidence for Lieb's conjecture for spin 2, so far a rigorous proof has been limited to certain symmetric configurations like equilateral triangles with centered fourth point ($\epsilon = \mu = \nu$ and $\alpha = \beta = \gamma$), and squares ($\alpha = \beta = \epsilon = \mu$ and $\gamma = \nu$). It is not hard to find values of the parameters that give values of S_W below the entropy for coherent states, but they do *not* correspond to any configuration of points on the sphere, so in contrast to spin 1 and spin 3/2 the constraint is now important. S_W is concave in each of the parameters ϵ, μ, ν, α, β, γ.

PROOF: The proof is analogous to the spin 1 and spin 3/2 cases but the geometry and algebra are considerably more complicated, so we will just give a sketch. Pick four points on the sphere, without loss of generality parametrized as $\omega_1^{(3)} = (\tilde{\theta}, 0)$, $\omega_2^{(3)} = (\theta, \phi)$, $\omega_3^{(3)} = (0, 0)$, and $\omega_4^{(3)} = (\bar{\theta}, \bar{\phi})$. Corresponding spin $\frac{1}{2}$ states are $|\omega_1^{(3)}\rangle$, $|\omega_2^{(3)}\rangle$, $|\omega_3^{(3)}\rangle$, as given in (34), (35), (36), and

$$|\omega_4^{(3)}\rangle = (1-\gamma)^{\frac{1}{2}} e^{\frac{-i\bar{\phi}}{2}} |\uparrow\rangle + \gamma^{\frac{1}{2}} e^{\frac{i\bar{\phi}}{2}} |\downarrow\rangle. \tag{50}$$

Up to normalization, the state of interest is

$$|\tilde{\psi}^{(3)}\rangle = P_{j=2} |\omega_1^{(3)} \otimes \omega_2^{(3)} \otimes \omega_3^{(3)} \otimes \omega_4^{(3)}\rangle. \tag{51}$$

In the computation of $|\tilde{\psi}_m^{(3)}|^2$ we encounter again the angle ϕ, compare (33),and two new angles $\bar{\phi}$ and $\bar{\phi} - \phi$. Luckily both can again be expressed as angles between planes that intersect the circle's center and we have

$$\begin{aligned} 2\cos\bar{\phi}\sqrt{\mu\gamma(1-\mu)(1-\gamma)} &= \mu + \gamma - \alpha - 2\mu\gamma, && (52)\\ 2\cos(\bar{\phi} - \phi)\sqrt{\epsilon\gamma(1-\epsilon)(1-\gamma)} &= \gamma + \epsilon - \beta - 2\gamma\epsilon, && (53) \end{aligned}$$

and find $1/c = \sum_m |\tilde{\psi}_m^{(3)}|^2$ as given in (45). By permuting the parameters ϵ, μ, ν, α, β, γ appropriately we can derive expressions for the remaining $|\tilde{\psi}_m^{(i)}|^2$'s and then compute S_W (44) with the help of (19). ∎

4 Higher spin

The construction outlined in the proof of corollary 3.5 can in principle also be applied to states of higher spin, but the expressions pretty quickly become quite unwieldy. It is, however, possible to use theorem 3.2 to show that the entropy is extremal for coherent states:

COROLLARY 4.1 (spin j) *Consider the state of spin j characterized by $2j-1$ coinciding points on the sphere and a $2j$'th point, a small (chordal) distance*

$\epsilon^{\frac{1}{2}}$ *away from them. The Wehrl entropy of this small deviation from a coherent state, up to third order in* ϵ*, is*

$$S_W(\epsilon) = \frac{2j}{2j+1} + \frac{c}{8j^2}\epsilon^2 \quad + \mathcal{O}[\epsilon^4], \tag{54}$$

with

$$\frac{1}{c} = 1 - \frac{2j-1}{2j}\epsilon \quad \textit{(exact)}. \tag{55}$$

A generalized version of Lieb's conjecture, analogous to (3), is [6]

CONJECTURE 4.2 *Let* $|\psi\rangle$ *be a normalized state of spin* j*, then*

$$(2js+1)\int \frac{d\Omega}{4\pi}\,|\langle\Omega|\psi\rangle|^{2s} \leq 1, \quad s > 1, \tag{56}$$

with equality if and only if $|\psi\rangle$ *is a coherent state.*

Remark: This conjecture is equivalent to the "quotient of two Hölder inequalities" (4). The original conjecture 2.1 follows from it in the limit $s \to 1$. For $s = 1$ we simply get the norm of the spin j state $|\psi\rangle$,

$$(2j+1)\int \frac{d\Omega}{4\pi}\,|\langle\Omega|\psi\rangle_j|^2 = |P_j|\psi\rangle|^2, \tag{57}$$

where P_j is the projector onto spin j. We have numerical evidence for low spin that an analog of conjecture 4.2 holds in fact for a much larger class of convex functions than x^s or $x \ln x$.

For $s \in \mathbb{N}$ there is a surprisingly simple group theoretic argument based on (57):

THEOREM 4.3 *Conjecture 4.2 holds for* $s \in \mathbb{N}$.

Remark: For spin 1 and spin 3/2 (at $s = 2$) this was first shown by Wolfgang Spitzer by direct computation of the integral.

PROOF: Let us consider $s = 2$, $|\psi\rangle \in [j]$ with $||\psi\rangle|^2 = 1$, rewrite (56) as follows and use (57)

$$\begin{aligned} &(2j \cdot 2 + 1) \int \frac{d\Omega}{4\pi} \, |\langle \Omega | \psi \rangle|^{2 \cdot 2} \\ &\qquad = (2(2j) + 1) \int \frac{d\Omega}{4\pi} \, |\langle \Omega \otimes \Omega | \psi \otimes \psi \rangle|^2 = |P_{2j} |\psi \otimes \psi\rangle|^2. \end{aligned} \tag{58}$$

But $|\psi\rangle \otimes |\psi\rangle \in [j] \otimes [j] = [2j] \oplus [2j-1] \oplus \ldots \oplus [0]$, so $|P_{2j}|\psi \otimes \psi\rangle|^2 < ||\psi \otimes \psi\rangle|^2 = 1$ unless $|\psi\rangle$ is a coherent state, in which case $|\psi\rangle \otimes |\psi\rangle \in [2j]$ and we have equality. The proof for all other $s \in \mathbb{N}$ is completely analogous. ■

It seems that there should also be a similar group theoretic proof for all real, positive s related to (infinite dimensional) spin js representations of su(2) (more precisely: sl(2)). There has been some progress and it is now clear that there will not be an argument as simple as the one given above [5]. Coherent states of the form discussed in [15] (for the hydrogen atom) could be of importance here, since they easily generalize to non-integer 'spin'.

Theorem 4.3 provides a quick, crude, lower limit on the entropy:

COROLLARY 4.4 *For states of spin j*

$$S_W(|\psi\rangle\langle\psi|) \geq \ln \frac{4j+1}{2j+1} > 0. \tag{59}$$

PROOF: This follows from Jensen's inequality and concavity of $\ln x$:

$$\begin{aligned} S_W(|\psi\rangle\langle\psi|) &= -(2j+1) \int \frac{d\Omega}{4\pi} \, |\langle \Omega | \psi \rangle|^2 \ln |\langle \Omega | \psi \rangle|^2 \\ &\geq -\ln \left((2j+1) \int \frac{d\Omega}{4\pi} \, |\langle \Omega | \psi \rangle|^{2 \cdot 2} \right) \\ &\geq -\ln \frac{2j+1}{4j+1}. \end{aligned} \tag{60}$$

In the last step we have used theorem 4.3. ■

We hope to have provided enough evidence to convince the reader that it is reasonable to expect that Lieb's conjecture is indeed true for all spin. All cases listed in Lieb's original article, 1/2, 1, 3/2, are now settled – it would be nice if someone could take care of the remaining "dot, dot, dot" …

Acknowledgments

I would like to thank Elliott Lieb for many discussions, constant support, encouragement, and for reading the manuscript. Much of the early work was done in collaboration with Wolfgang Spitzer. Theorem 4.3 for spin 1 and spin 3/2 at $s = 2$ is due to him and his input was crucial in eliminating many other plausible approaches. I would like to thank him for many discussions and excellent team work. I would like to thank Branislav Jurčo for joint work on the group theoretic aspects of the problem and stimulating discussions about coherent states. It is a pleasure to thank Rafael Benguria, Almut Burchard, Dirk Hundertmark, Larry Thomas, and Pavel Winternitz for many valuable discussions. Financial support by the Max Kade Foundation is gratefully acknowledged.

References

[1] A. Anderson, J.J. Halliwell, *Information-theoretic measure of uncertainty due to quantum and thermal fluctuations*, Phys. Rev. D **48** (1993) 2753–2765

[2] W. Beckner, *Inequalities in Fourier analysis*, Ann. of Math. **102** (1975) 159–182

[3] H.J. Brascamp, E.H. Lieb, *Best constants in Young's inequality, its converse, and its generalization*, Adv. in Math. **20** (1976) 151–173

[4] M. Grabowski, *Wehrl-Lieb's inequality for entropy and the uncertainty relation*, Rep. Math. Phys. **20** (1984) 153–155

[5] joint work with B. Jurčo

[6] E.H. Lieb, *Proof of an entropy conjecture of Wehrl*, Commun. Math. Phys. **62** (1978) 35–41

[7] E.H. Lieb, *Coherent states as a tool for obtaining rigorous bounds*, In: D.H. Feng, J. Klauder (eds.), Coherent states: past, present, and future. Proceedings, Oak Ridge, pp. 267 – 278. World Scientific 1994

[8] E.H. Lieb, *Some convexity and subadditivity properties of entropy*, Bull. Amer. Math. Soc. **81** (1975) 1–13

[9] E.H. Lieb, *Gaussian kernels have only Gaussian maximizers*, Invent. Math. **102** (1990) 179–208

[10] E.H. Lieb, *Integral bounds for radar ambiguity functions and Wigner distributions*, J. Math. Phys. **31**, (1990) 594–599

[11] E.H. Lieb, M. Loss, Analysis, Graduate Studies in Mathematics, 14. American Mathematical Society, Providence, RI, 1997

[12] J. Peřina, Z. Hradil, B. Jurčo, Quantum optics and fundamentals of physics, Fundamental Theories of Physics, 63. Kluwer Academic Publishers Group, Dordrecht 1994

[13] J.M. Radcliffe, *Some properties of coherent spin states*, J. Phys. A **4**, (1971) 313–323

[14] joint work with W. Spitzer

[15] L.E. Thomas, C. Villegas-Blas, *Asymptotic of Rydberg states for the hydrogen atom*, Commun. Math. Phys. **187** (1997) 623–645

[16] A. Wehrl, *On the relation between classical and quantum mechanical entropy*, Rept. Math. Phys. **16** (1979) 353–358

[17] W.-M. Zhang, D.H. Feng, R. Gilmore, *Coherent states: Theory and some applications*, Rev. mod. Phys. **62** (1990) 867–927